



As folhas do sanguinho-de-água servem de alimento às lagartas da Borboleta-limão e Spini.

The caterpillars of Brimstone and Ilex Hairstreak feed on alder buckthorn's leaves.

© Frank Pennelkamp

SANGUINHO-DE-ÁGUA | Alder buckthorn
Frangula alnus
Eudicotiledonea / Rhamnaceae



BORBOLETA-LIMÃO | Brimstone
Gonepteryx rhamni
Insecta / Lepidoptera / Pieridae

© Albano Soares



SPINI | Ilex Hairstreak
Satyrium spini
Insecta / Lepidoptera / Lycaenidae

© Frank Pennelkamp



Esta borboleta coloca os seus ovos em folhas de carvalho-negral ou de silvas, assim, quando as lagartas nascem têm alimento à disposição.

This butterfly lays its eggs on the leaves on pyrenean oaks or elmleaf blackberries, so that caterpillars are born on a good source of food.

© Albano Soares

AZUL-CELESTE | Holly Blue
Celastrina argiolus
Insecta / Lepidoptera / Lycaenidae



CARVALHO-NEGRAL | Pyrenean oak
Quercus pyrenaica
Eudicotiledonea / Fagaceae

© Frank Pennelkamp



SILVA | Elmleaf blackberry
Rubus ulmifolius
Eudicotiledonea / Rosaceae

© Frank Pennelkamp